Chemistry for Medicine

Name: MODEL ANSWERS	ID Number:
---------------------	------------

Time: 2 hours

Useful constants: $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

1 amu = 1.6605×10^{-24} g

1 atm = 760 mmHg = 760 torr

1 H 1.008																	2 He 4.003
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
55 Cs 132.9	56 Ba 137.3	57 La* 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.9	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226	89 Ac [†] (227)								,							

QUESTION	SCORE	MAXIMUM MARKS
1		
2		
TOTAL		

QUESTION 1

(a) A medical student was asked to determine the molarity of an aqueous solution of citric acid in the laboratory using a titration with sodium hydroxide. The molarity of the citric acid solution was known accurately to be 0.0560 mol/L.

The student performed the experiment four times and obtained the following results:

0.0533 mol/L

0.0530 mol/L

0.0534 mol/L

0.0531 mol/L

(i) In the laboratory what glassware do we use to measure volumes of liquids in general?

volumetric flask

(ii) The aqueous solution of citric acid was prepared from a crystalline solid of citric acid.

What are the different methods used to measure mass in the laboratory?

Weighing by difference

Name the two types of balances used to measure mass in the laboratory.

analytical balance top-loading balance

(iii) Mass is an extensive property. What does this mean?

It means that it depends on the amount of the substance considered

Give two examples of an intensive property. density and temperature

(iv) **Discuss** the results of the determination of the molarity of citric acid above.

Since the measured values of the molarity are close to each other the measurements are precise or reproducible. The average value of the measurements is as follows:

(0.0533 mol + 0.0530 mol + 0.0534 mol + 0.0531 mol + 0.0

= 0.0532 mol/L

The % error = $(0.0560 - 0.0532) \times 100\% = \frac{0.0028}{0.0560} \times 100\%$

The average value differs slightly from the actual (true) molarity. Thus the measurements are not very accurate, i.e. they are slightly accurate. (v) Name the type of reaction that was used in the analysis of the citric acid solution.

acid-base reaction or neutralisation reaction

- (b) Answer the following questions about hemoglobin.
- (i) What is hemoglobin? An iron-binding protein that transports oxygen from the lungs to the body tissues via the blood circulatory system.
 - (ii) How many iron atoms are found in one molecule of hemoglobin?

(iii) Hemoglobin is 0.3437% by mass of iron. Calculate the molecular mass of hemoglobin.

$$0.3437\% = \frac{4(55.85)}{M} \times 100\%$$
 $M = 6.500 \times 10^4 \text{ amu} = \text{molecular mass}$

(iv) An average adult has 5.0 L of blood. Every milliliter (mL) of blood contains 5.0×10^9 red blood cells. Every red blood cell has an average of 2.8×10^8 molecules of hemoglobin.

Calculate the total mass of hemoglobin (in grams) in an adult.

5.0 L
$$\times \frac{1 \text{mL}}{10^{3} \text{L}} \times \frac{5.0 \times 10^{9} \text{ RBC}}{1 \text{ mL}} \times \frac{2.8 \times 10^{8} \text{ molecules of Hb}}{1 \text{ RBC}}$$

= $7.0 \times 10^{21} \text{ Hb molecules}$

= $\frac{10^{21} \text{ Hb molecules}}{1 \text{ Hb}} = \frac{10^{21} \text{ Hb}}{1 \text{ Hb}} = \frac{10^{21} \text{ Hb}$

Determine the total mass of iron in the hemoglobin of an adult.

Total mass of iron in
$$Hb = 0.3437\% \times 750g$$

= $2.6g$

The mass of iron in hemoglobin is 65% of the total amount of iron in an adult. Calculate the total mass of iron in an adult.

$$65\% = \frac{m_{Fe(Hb)} \times 100\%}{m_{Fe(body)}} \times 100\%$$

$$65\% = \frac{2.69}{M_{Fe(body)}} \times 100\% : m_{Fe(body)} = 4.09$$

(c) An unknown organic compound contains carbon, hydrogen and oxygen.

Complete combustion of 1.2010 g of this compound gives 1.7604 g of carbon dioxide. Complete combustion of 36.00 mg of this compound produces 7.220 × 10²⁰ molecules of water.

(i) Determine the empirical formula of the unknown organic compound.

molecular formula of this compound?

One molecule of CxHyOz contains 12 atoms = 4 atoms : Molecular formula = Cx H40z But since the empirical formula is CH2O, the molecular formula is C2H402

- (d) Write reaction equations for each of the following:
- (i) The combustion of ethanol

(ii) When aluminium is added to hydrobromic acid hydrogen gas is given off.

(iii) The heating of a hydrated alkaline earth metal halide produces an anhydrous salt. (Alkaline earth metal = **M**, halogen = **X**)

$$M \times_2 \cdot n +_2 \circ (s) \xrightarrow{\Delta} M \times_2 (s) + n +_2 \circ (g)$$

(iv) Mixing an aqueous solution of mercury(I) nitrate with an aqueous solution of ammonium phosphate produces a precipitate. (All salts of ammonium are soluble in water).

(v) Sublimation of carbon dioxide.

$$CO_2(S) \xrightarrow{\Delta} CO_2(G)$$

(vi) The oxidation of iron by oxygen gas produces a solid of iron(III) oxide as the only product.

(vii) The reaction of formic acid with calcium hydrogen carbonate produces a colourless gas.

(viii) The reaction of tetraphosphorus decoxide with water forms phosphoric acid as the only product.

(ix) The hydronium ion and the hydroxide ion react in aqueous solution to produce water.

(x) The thermal dissociation of a molecule of iodine.

$$T_2(s) \xrightarrow{\Delta} 2I(g)$$

QUESTION 2

(a) Perform the following mathematical operations. Show all your calculations.

(i)
$$20 \circ C = 2.9 \times 10^2 \times 20 + 273.15 \Rightarrow 29. \times 10^{\frac{1}{2}} \times 10^$$

(ii) 646 torr =
$$\frac{0.850}{1000}$$
 atm
646 torr $\times \frac{1000}{7600} = 0.850$ atm

(iii)
$$9500 + 4500 = 1.40 \times 10^{4}$$

$$+ 4.5 \times 10^{3}$$

$$+ 4.5 \times 10^{3}$$

$$+ 4.0 \times 10^{4}$$

$$\Rightarrow 1.40 \times 10^{4}$$

(iv)
$$0.03036 \text{ cm} \div 50.0 \text{ cm} = 6.07 \times 10^{-4}$$

$$\frac{0.03036 \text{ cm}}{y} = 6.07 \times 10^{-4}$$

$$\therefore y = \frac{0.03036 \text{ cm}}{6.07 \times 10^{-4}} = 50.0 \text{ cm}$$

(v)
$$0.039 \, dm^3 + 60.8 \, mL = 100 \cdot mL \, or \, 1.00 \times 10^2 ... \rightarrow 39 \, mL$$

 $0.039 \, dm^3 \times \frac{L}{dm^3} \times \frac{mL}{10^{-3}L} = 39 \, mL \longrightarrow 99 \cdot 8 \, mL$

$$9.9862 \times 10^{2}$$
(vi) 9.9862×10^{5} mg/dm³ = $\frac{9.9862 \times 10^{2}}{9.9862 \times 10^{5}}$ µg/µL
$$9.9862 \times 10^{5}$$
 mg $\times \frac{10^{3}}{10^{6}} \times \frac{10^{6}}{10^{6}} \times \frac{10^$

(b) Chlorine has two isotopes. One of these isotopes is ³⁵Cl with an atomic mass of 34.93 amu and a natural abundance of 75.53%.

Calculate the mass of the other isotope of chlorine in grams.

35C1 34.93 amu 75.53%
×C1 m?
$$100\%-75.53\% = 24.47\%$$

$$\frac{75.53}{100}(34.93) + \frac{24.47}{100}(m) = 35.45$$

$$2638 + 24.47 m = 3545$$

$$24.47 m = 907$$

$$m = 907 = 37.1$$

$$\therefore \text{ atomic mass of Second isotope of } C1 = 37.1 \text{ amu}$$

$$\text{mass in grams} = 37.1 \text{ amu} \times \frac{1.6605 \times 10^{-24}}{24.47}g$$

$$= 6.16 \times 10^{-23}g$$

(c) Aluminium reacts with liquid bromine (Br₂) to produce solid aluminium bromide.

In an experiment, 40.75 g bromine was reacted with 5.56 g aluminium and 38.54 g aluminium bromide was obtained.

Calculate the percentage yield of aluminium bromide.

$$2AI(s) + 3Br_2(s) \longrightarrow 2AIBr_3(s)$$

$$\frac{5.56}{2}g + \frac{40.75}{9}g = \frac{40.75}{159.8}g/mol + \frac{1}{159.8}g/mol + \frac{$$